

# The effect of time-trial duration on aerodynamic drag

A. Bouillod<sup>1,2,3</sup>, E. Brunet<sup>2</sup>, G. Soto-Romero<sup>3,4</sup>, L. Garbellotto<sup>3</sup>, G. Millour<sup>5</sup> & F. Grappe<sup>1,6</sup>

<sup>1</sup> EA4660, C3S Health - Sport Department, Sports University, Besancon, France

<sup>2</sup> French Cycling Federation, Saint Quentin en Yvelines, France

<sup>3</sup> LAAS-CNRS, Université de Toulouse, CNRS, Toulouse, France

<sup>4</sup> ISIFC, Université de Franche-Comté, France

<sup>5</sup> Laboratoire M2S, Université Rennes 2 – ENS Rennes, Rennes, France

<sup>6</sup> Professional Cycling Team FDJ, Mousy le Vieux, France

**Keywords:** cycling position, aerodynamic drag, power output, time-trial.

**Purpose:** Aerodynamic drag is the main resistance (80-90 %) among the total resistive forces ( $R_T$ , N) opposing motion on level ground in cycling (Debraux et al. 2011). To reduce air resistance, cyclists adopt a characteristic time trial (TT) position on the bicycle to decrease the frontal area. The bike frame and the components only accounts for about 30 % of the total drag of the cyclist-bicycle system. Thus, the cyclist position has a great importance in the performance on flat terrain to overcome at the maximum the air resistance (Oggiano et al. 2008). Many research have studied the effect of the aerodynamic position or equipment's on the air resistance but to the best of our knowledge, no study has yet reported the effect of the duration of the exercise on aerodynamic drag. The aim of this study was to analyse the effect of time-trial duration on aerodynamic drag. We hypothesized that the more the duration of exercise is long the more the fatigue increases involving an alteration of the effective frontal area ( $AC_d$ , m<sup>2</sup>) and accordingly a decrease of performance.


**Methods:** 9 elite road cyclists performed all testing session with their personal TT bike on a 200 m covered velodrome (Bourges, France). The bicycle was fitted with a rear wheel composed of a Powertap G3 hub (CycleOps, Madison, USA) for the measurement of speed ( $V$ , m.s<sup>-1</sup>) and power output (PO, W). Firstly, the cyclists performed a discontinuous incremental exercise at different  $V$  to determine the rolling resistance coefficient (Cr) from the  $R_T$ - $V^2$  linear regression method (Grappe et al. 1997). Then, the Cr was fixed to compute the  $AC_d$  during three TTs of different durations according to the cyclists' categories (table 1). The effects of TT duration and time on  $AC_d$ ,  $V$  and PO were tested using a two-way analysis of variance (ANOVA).

**Table 1:** TT durations (min) according to the categories.

Categories	TT1	TT2	TT3
U17 Men / U19 Women	3	8	15
U19 Men / U23 Women	4	10	20
U23 + Elite Men / Elite Women	5	15	30

**Results:** There was no significant effect of TT duration and time on  $AC_d$ . The largest decrease in  $AC_d$  was during TT1 (-2.9 %). PO and  $V$  were significantly influenced by TT duration ( $p < 0.001$ ). The mean  $V$  was significantly different ( $p < 0.001$ ) in all TT durations whereas the mean PO was significantly higher ( $p < 0.001$ ) in TT1 compared to TT2 (+20.5 %) and TT3

(+28.2 %). A significant correlation was measured between  $V$  and the  $PO/AC_d$  ratio in TT1 ( $r = 0.94$ ,  $p < 0.001$ ), TT2 ( $r = 0.95$ ,  $p < 0.001$ ) and TT3 ( $r = 0.97$ ,  $p < 0.001$ ).



**Figure 1.** Mean changes (%) in  $AC_d$  and PO during the three time-based cycling time trials.  
*\*The standard deviations don't appear to simplify the figure.*

**Conclusions:** The main results show that the mean  $AC_d$  decreased non-significantly over time for all the TT durations suggesting that the position of the cyclists was not altered during the three events. According to unpublished studies conducted in our lab, wind tunnel sessions have demonstrated that the  $AC_d$  decreases by a mean of  $0.5\%$  per  $1\text{ km}\cdot\text{h}^{-1}$  for speeds between  $40$  and  $60\text{ km}\cdot\text{h}^{-1}$ . Even more, by applying a correction coefficient on  $AC_d$ , the lowest value ( $-2.9\%$ ) will be reduced to  $-2.3\%$  which reinforces the fact that  $AC_d$  remains relatively stable during the effort. The strong correlations between  $V$  and the  $PO/AC_d$  in the three TTs demonstrate that the more the  $PO/AC_d$  was high, the more the  $V$  was high and the performance was improved (Peterman et al. 2015). Thus, the balance between PO and  $AC_d$  is a parameter that must be optimised taking into account both the training process and aerodynamic resistance. To be efficient in a TT, the cyclist will find the better aero-position on his bike being able to develop the most important level of PO.

## References:

1. Debraux P, Grappe F, Manolova AV, Bertucci W (2011) Aerodynamic drag in cycling: methods of assessment. *Sports biomechanics / International Society of Biomechanics in Sports* 10: 197-218
2. Grappe F, Candau R, Belli A, Rouillon JD (1997) Aerodynamic drag in field cycling with special reference to the Obree's position. *Ergonomics* 40: 1299-1311
3. Oggiano L, Leirdal S, Saetran L, Ettema G (2008) Aerodynamic optimization and energy saving of cycling postures for international elite level cyclists. ISEA Conference, Biarritz
4. Peterman JE, Lim AC, Ignatz RI, Edwards AG, Byrnes WC (2015) Field-measured drag area is a key correlate of level cycling time trial performance. *PeerJ* 3: e1144